

Locality and reality revisited

Adrian Kent^{1,2}

¹ *Hewlett-Packard Laboratories, Filton Road,
Stoke Gifford, Bristol BS34 8QZ, U.K.*

(February 2002)

Bell gave the now standard definition of a local hidden variable theory and showed that such theories cannot reproduce the predictions of quantum mechanics without violating his “free will” criterion: experimenters’ measurement choices can be assumed to be uncorrelated with properties of the measured system prior to measurement.

An alternative is considered here: a probabilistic theory of hidden variables underlying quantum mechanics could be statistically local, in the sense that it supplies global configuration probabilities which are defined by expressions involving only local terms. This allows Bell correlations without relying on *either* a conspiracy theory in which prior common causes correlate the system state with experimenters’ choices *or* a reverse causation principle in which experimenters’ choices affect the earlier system states. In particular, there is no violation of the free will criterion. It gives a different perspective on Bell correlations, in which the puzzle is not that apparently non-local correlations should emerge from rules involving local quantities, but rather that we do not see more general non-local correlations that allow paradox-immune forms of superluminal signalling.

² On leave from DAMTP, University of Cambridge, Silver Street, Cambridge CB3 9EW, U.K.

I. A LAMENT

Many important concepts in quantum foundations are misleadingly named:

- The *measurement problem* is not the problem of understanding precisely what a measurement is — something which needn’t be, and probably isn’t, specified in the fundamental theory of nature — but the problem of finding a precise mathematical or physical characterisation of the sample space on which quantum probabilities are defined.
- *Hidden variables* are hidden in the sense that the quantum formalism conceals them, but, as Bell emphasized, certainly should not be supposed to be hidden from us, since they are meant to define the reality we experience. And Bell’s own preferred term, *beable*, has a metaphysically confusing connotation of a not necessarily actualised potential for existence. This would seem to make sense for the values of suitable variables — only one value is realised out of many possibilities — but not for the variables themselves: nonetheless, Bell intended the latter. Better terms for quantities parametrising the sample space which solves the measurement problem would be *existents* (Shimony’s term [2]) or perhaps *real variables*.
- Bell’s *free will criterion* [3] conflicts with some widely held views of what it means to be able to incorporate free will into a physical theory — and Bell’s response to critics explains that he did not actually intend to express a view on the problem of free will. Free will, as Bell used it, is no more than a convenient shorthand term. A theory satisfies the free will criterion if it allows us to justify treating the measurement settings as independent variables to those characterising the state of the microscopic system (typically, a pair of separated entangled particles) being measured. This need not mean that the theory admits your, or anybody’s, favourite notion of free will. Conversely, as Price emphasizes [4], a theory which fails to satisfy the free will criterion may nonetheless be compatible with sensible definitions of free will.
- Finally, *quantum non-locality* is a confusingly oxymoronic phrase. Quantum field theories — the best version of quantum theory we have — do satisfy a precise locality condition: operators with support in space-like separated regions commute. What the term is (or should be) meant to signify is that quantum theory is not locally causal under Bell’s definitions — but *quantum non-(local causality)* is too clumsy.

What to do? Using non-standard terms, or inventing new ones, makes life difficult for readers. So I will use the standard terms, with these criticisms understood.

II. SOME MOTIVATION

It seems to me that all our difficulties in quantum foundations stem first from the measurement problem, and then from the fact that the obvious ways of tackling the measurement problem are hard to reconcile with special relativity and/or reality.

Probably this is because quantum theory is wrong, and the insolubility of the measurement problem is a symptom. Just possibly, another interpretation of quantum theory might do the trick.

Either way, there's a case for carefully re-examining the conventional lines of argument, looking for gaps and trying to find scope for something new. Maybe there's also a case for giving up and waiting for experiment to give us a clue, but I'll resist that temptation here. Instead, I want to describe a different way of thinking about hidden variables, which at least sheds a different light on Bell correlations.

III. LATTICE MODELS IN STATISTICAL PHYSICS

To statistical physicists, lattice models are mathematical idealizations. Many quantities of interest in statistical physics are relatively insensitive to the details of the microphysics, and many distinct systems and models may belong to the same universality class, with essentially the same properties. So, if a well chosen lattice model reproduces the bulk properties and correlation functions observed in an experimental sample, this need not imply that the model even approximately describes the system's microphysics.

I want here to use some standard definitions of lattice statistical physics in an unfamiliar way. First, I note that lattice-like models make sense as fundamental theories in their own right, rather than as phenomenological models attempting to capture some of the properties of a more fundamental theory. Then I want to consider the possibility that something rather like a lattice model could be a fundamental theory of microphysics, giving a hidden variable theory explanation of Bell correlations.

Consider the two-dimensional Ising model on a square lattice (Figure 1). A "spin" variable $s_i \in \{-1, 1\}$ is attached to each site i . The Hamiltonian

$$H = \sum_{(i,j) \text{ nearest neighbours}} (1 - s_i s_j),$$

includes only interactions between horizontally or vertically adjacent spins. For any possible configuration of spins $S = \{s_i : i \text{ a lattice site}\}$, we have a *probability weight*

$$p(S) = \exp(-CH),$$

where C is a constant (at fixed temperature). The idealised system represented by the Ising model will always be found in some legitimate spin configuration, and the probability of finding it in the configuration S is

$$\text{Prob}(S) = \frac{p(S)}{\sum_{\text{all configurations } S'} p(S')}.$$

Note that the model is thus *statistically local*, in the sense that if two spin configurations S and S' differ only on a region R , the ratio

$$\frac{\text{Prob}(S)}{\text{Prob}(S')}$$

can be calculated purely from the values of the s_i and s'_i within the region R , together with those of their nearest neighbours. That is, if S and S' differ only locally, the ratio of their probabilities can be calculated from local quantities.




Figure 1: An Ising model spin configuration

Now, the main point. *The Ising model, as just described, could be interpreted as a fundamental theory.* Imagine a world in which, say, a particular type of crystal has the property that its surfaces are perfect square arrays of atoms, in which each atom has an asymmetric feature which define an orientation, and in which the orientation is always perpendicular to the surface, either pointing outward (+1) or inward (−1). Imagine further that the orientations are stable — we cannot change them — and can be examined by powerful microscopes, and that a plentiful supply of crystals is available.

Then it makes sense, as a hypothesis, to suppose that the atomic orientations are randomly configured, with probabilities defined by the Ising model (on the finite lattice defined by the crystal surface). The theory can be tested: examining large numbers of crystals will reveal whether or not the data are statistically consistent with the theory. And it need not be a phenomenological theory; it could be fundamental. It could simply be that nature has fixed these unchanging orientation configurations, not through some underlying dynamics whose end result is described by the Ising model, but from the beginning, by a random choice which (like the randomness of quantum measurements, according to the orthodox view) has no underlying explanation.

Obviously, this imaginary Ising-world is not much like ours, in which crystals form and decompose, atoms don't behave classically, their configurations and spin states aren't fixed, and everything is (as far as we presently understand) ultimately described by quantum dynamics. But conceptually the Ising theory of crystals in Ising-world makes sense. It is testable in much the same way as quantum theory is, and it includes a simplifying principle — statistical locality — which makes it much easier to test: it suffices to look at orientation configurations in local regions of crystal surfaces. And, given our other assumptions about Ising-world, there is no compelling reason — nothing analogous to the measurement problem — to think Ising theory must be incomplete.

IV. MINKOWSKI SPACE TRAJECTORY MODELS

Let us suppose that underlying quantum theory there is some theory in which real particles follow definite trajectories through spacetime. To simplify things, take space-time to be two-dimensional, with a discrete lattice structure at

microscopic scales (Figure 2). The lattice edges are lightlike, and so the discussion is restricted to massless particles, which travel along the lattice edges. Also, for the purposes of this discussion, macroscopic experimental apparatus will be treated as qualitatively distinct from microscopic particles, so that we are allowed to consider apparatus set up somewhere in spacetime without having to describe it as a collection of microscopic particles, each following their own trajectories through the lattice.

All of these simplifications are meant to be provisional rather than fundamental. The idea is to set up the standard discussion of Bell correlations, and then find a new way of looking at them, without obscuring things with too many technicalities.




Figure 2: Two-dimensional Minkowski space trajectory lattice

Figure 3 shows a standard Bell experiment on a pair of entangled massless particles, “photons”, with one extra feature — lightlike trajectories along the lattice edges have been included. Though this looks like a standard beam path picture, the trajectories here are hidden variables: the picture is meant to be interpreted as saying that the photons *really* did travel along the paths depicted — something we will eventually have to show is consistent with the model we produce. The space-time trajectories of the classical measuring apparatuses won’t affect the discussion, and are omitted.




Figure 3: A Bell experiment

To further simplify things — since we cannot use photon polarizations in two dimensions — let us assume that the measurements are perfect von Neumann measurements of an internal degree of freedom, “spin”, with the photons initially in the singlet state of a pair of spin 1/2 particles.¹ According to quantum theory, these von Neumann measurements do not affect the form of the photon wave packet, assuming (let’s) that the spatial and spin degrees of freedom are unentangled. It seems reasonable to postulate that, in the hidden variable model we are building, the photon trajectories will also be unaffected by such measurements. For instance, after a measurement of spin in the measurement basis defined by vector $\mathbf{a}$, the apparatus records the answer $\mathbf{a}$ or $-\mathbf{a}$, the photon emerges in the corresponding spin eigenstate, and its trajectory continues unaltered.

The aim now is to reproduce Bell correlations by a statistically local lattice-like model involving photon trajectories, while respecting Bell’s free will criterion. We will want to attach further hidden variable degrees of freedom to the photon trajectories, so that the spin measurements become relevant to the hidden variable model. The free will criterion requires that no hidden variables at points before the measurements should be correlated with the measurement choices. This, together with locality, means that the photons’ states before the measurements cannot explain the correlations: we need rules involving events *after* the measurements.

We can ensure a relevant post-measurement event if the photons are forced to meet again, and *that* can be done, without violating statistical locality, as follows. First, we give a small but non-zero probability weight to trajectory segments in which a photon’s direction switches: this allows the two photons to meet after the measurements. Second, we choose rules that assign probability zero to the event that the photons propagate forever, and non-zero probability weight to the event that two photon trajectories annihilate when they meet. Together these ensure that photons *must* eventually undergo pair annihilation. The first of these can be arranged by assigning probability weight of $(1 - \epsilon)$ to a vertex in which the photon continues in the same direction, and weight ϵ to a vertex in which it switches (Figure 4); here $\epsilon > 0$ is chosen to be small enough that $(1 - \epsilon)^N \approx 1$, where N is the number of lattice links a photon would traverse in the largest scale Bell experiment carried out to date.

¹The scare quotes around “photon” and “spin” are left implicit from now on.




Figure 4: Probability Weights for Free Propagation

In choosing rules for a pair annihilation, we need to make sure that the originally entangled photons meet and mutually annihilate each other, and not other photons. This we do by introducing a hidden variable label which only the entangled pair share, and allowing annihilation only between pairs which share the same label. Finally, we enforce the correct correlations by fiat, by adding a further label which records a photon's measurement outcome, and by making the probability weight for the annihilation event depend appropriately on these labels (Figure 5).




Figure 5: Probability Weight for Recombining
Bell Particles after Measurements

To summarise: we've arranged rules so that the entangled photons from pair i do the following: (i) with high probability, propagate linearly from source to detectors, (ii) after measurement, each eventually undergo one or more changes of direction, (iii) with probability one, eventually meet up and annihilate. (Figure 6)

The trajectory from A to V carries hidden variables i (pair label) and a (measurement result)
The trajectory from B to V carries hidden variables i (pair label) and b (measurement result)
The trajectories from S to A and B carry hidden variable i (pair label)




Figure 6: a typical trajectory description of a Bell experiment in the model

V. DISCUSSION

All we've done, of course, is rewrite the standard quantum correlations into ad hoc rules, in such a way that we can formally attach hidden variables characterising the measurement results to the photon paths after measurement, and formally account for the correlations as deriving from a rule attached to a pair annihilation event that must take place in the far future. What was the point of that?

The point, as I see it, is to underline how difficult it is to draw absolutely watertight conclusions about the form of realist theories underlying quantum theory, even if we accept that such theories should respect special relativity and should incorporate some form of locality, and even after decades of analysis of the implications of Bell experiments. To recapitulate:

We can agree with Bell that any theory which postulates common causes in the past which systematically affect experimenter and system, in such a way as to always produce Bell correlations, is likely to be unbelievably conspiratorial. But, as Price has so carefully and persuasively argued [4], this leaves open the possibility of theories which invoke some form of advanced action principle, by which experimenters' choices affect the earlier state of the system. As Price notes, although such theories would violate Bell's free will criterion, they would not necessarily rule out free will *per se*, at least according to some reasonable definitions. And they would not contradict the notion of effective freedom which underlay Bell's definition, nor invoke the same type of implausible conspiracy as a common cause explanation.

Still, there is (at the very least) an aesthetic case for the free will criterion. But even if the criterion is accepted, the discussion isn't closed. The models described above respect the free will criterion and they invoke no complex common cause hypothesis. They incorporate a respectable form of locality. Yet they give — in the simple case considered — a consistent explanation of Bell experiments.

That said, there is, of course, much to be said against the sort of ideas outlined above. The rules proposed are entirely ad hoc, transparently constructed precisely in order to reproduce quantum correlations. They could be generalised to cover more complicated entanglement experiments, but only — it would seem — at the price of introducing even more ad hoc and unwieldy rules. They require, in particular, that every hidden particle trajectory carries a record of its past history — which particles it was originally entangled with, and which measurements were carried out on it — arbitrarily far into the future. It is hard to see, methodologically speaking, what the point of pursuing this idea would be, unless some independent principle could be found to explain why the annihilation vertex weights should take the form necessary to reproduce quantum correlations. And where could such a principle come from?

In fact, any type of correlation, including those which allow superluminal signalling, could consistently be attached to the annihilation vertex. Why should such correlations be excluded? If we were to take the basic idea of these models seriously, we would be led to ask why the non-local correlations we see in nature are so tame as to preclude superluminal signalling. That this would lead to paradoxes is not a valid answer. Consistent probabilistic rules for global configurations cannot produce a logical contradiction. Attempts to set up a paradoxical causal loop fail. This is essentially because the probability of a successful superluminal signal is — even in isolation — always at least slightly less than one, and the probability of a sequence of signals succeeding is *not* generally the product of the values of the individual probabilities of success if the relevant signalling devices were isolated [5].

Moreover, the model involved several breathtaking simplifications, which would need to be removed if the idea were to be taken seriously. (Can we eliminate the need for an underlying lattice, or ensure that the model has a Lorentz invariant continuum limit? What about rules for massive particles, and for other types of measurement? And how do we build a coherent model that describes macroscopic measuring devices in terms of microscopic particles?)

Nonetheless, despite all these flaws, I think there is some interest in this rewriting of the standard story. It seems to show that the conventional wisdom about Bell's "free will" criterion is not quite right: realist, local and Lorentz invariant theories can respect the criterion and yet reproduce quantum correlations. And it does so by identifying a class of realist models — probabilistic models in which the probability of a configuration depends on a product of local terms — which standard discussions of Bell's theorem neglect. There are, as just noted, many arguments against taking seriously the idea that nature is actually described by these models. But the arguments are different from the arguments against other unorthodox ideas about hidden variables and Bell correlations, and worth thinking through, just for that reason.

VI. ACKNOWLEDGEMENTS

I am very grateful to Tomasz Placek and Jeremy Butterfield for giving me the opportunity to present this work at a most stimulating and enjoyable meeting, and for patient editorial encouragement far beyond the call of duty. The work was supported in part by a Royal Society University Research Fellowship, PPARC, and the European collaboration EQUIP.

- [1] J.S. Bell, *The theory of local beables*, Epistemological Letters, March 1976, reprinted in *Dialectica* **39** 85-96 (1985).
- [2] A. Shimony, *Reply to Bell*, *Dialectica* **39** 107-110 (1985).
- [3] J.S. Bell, *Free variables and local causality*, Epistemological Letters, February 1977; reprinted in *Dialectica* **39** 103-106 (1985).
- [4] H. Price, *Time's Arrow and Archimedes' Point: New Directions for the Physics of Time*, Oxford University Press, Oxford (1996).
- [5] See e.g. A. Kent, *Causality in Time-Neutral Cosmologies*, *Phys. Rev. D* **59** 043505 1-5 (1998).